arXiv:1612.09495v1 [cs.IT] 30 Dec 2016

The (n, m, k, λ)-Strong External Difference Family with $m \geq 5$ Exists *

Jiejing Wen, Minghui Yang and Keqin Feng

Abstract The notion of strong external difference family (SEDF) in a finite abelian group $(G, +)$ is raised by M. B. Paterson and D. R. Stinson [5] in 2016 and motivated by its application in communication theory to construct R-optimal regular algebraic manipulation detection code. A series of (n, m, k, λ)-SEDF's have been constructed in [5, 4, 2, 1] with $m = 2$. In this note we present an example of (243, 11, 22, 20)-SEDF in finite field $\mathbb{F}_q$ ($q = 3^5 = 243$). This is an answer for the following problem raised in [5] and continuously asked in [4, 2, 1]: if there exists an (n, m, k, λ)-SEDF for $m \geq 5$.
Key Words strong external difference family, cyclotomic class, cyclotomic number, finite field, strong algebraic manipulation detection code.

1 Introduction

Let $(G, +)$ be a finite abelian group with $n = |G|$ elements. For subsets D_1 and D_2 of G ($|D_1|, |D_2| \geq 1$), we defined the following multiset

$$\Delta(D_1, D_2) = \{a_1 - a_2 : a_1 \in D_1, a_2 \in D_2\}$$

Definition Let $A_1, \ldots, A_m$ ($m \geq 2$) be subsets of G, $|A_i| = k$ ($1 \leq i \leq m$). The family $\{A_1, \ldots, A_m\}$ is called an (n, m, k, λ)-strong external difference family (SEDF) in G if for each i ($1 \leq i \leq m$),

$$\sum_{\substack{j=1 \\ j \neq i}}^{m} \Delta(D_i, D_j) = \lambda(G - \{0\}) \qquad (1)$$

where we use the notation in the group ring $\mathbb{Z}[G]$ to express the multisets in both sides of equality (1). Namely, this equality means that for each nonzero element g in G, the multiplicity of g in the multiset

$$\bigcup_{\substack{j=1 \\ j \neq i}}^{m} \Delta(D_i, D_j)$$

* The work of J. Wen and M. Yang is supported by the NSFC under Grant 61171082, 61379139. The work of K. Feng was supported by the NSFC under Grant 11471178, 11571107.
J. Wen is with the Chern Institute of Mathematics, Nankai University, Tianjin 300071, China(e-mail:jjwen@mail.nankai.edu.cn)
M. Yang is with the State Key Laboratory of Information Security, Institute of Information Engineering, Chinese Academy of Sciences, Beijing 100195, China(e-mail:yangminghui6688@163.com)
K. Feng is with the Department of Mathematical Sciences, Tsinghua University, Beijing, 100084, China(e-mail: kfeng@math.tsinghua.edu.cn)

is constant λ, and 0 is not in this multiset. It is easy to see that the equality (1) implies that $(m-1)k^2 = \lambda(n-1)$ and $A_1, \ldots, A_m$ should be pairwise disjoint.

Let $G = \{g_1, \ldots, g_n\}$. Then $\{\{g_1\}, \ldots, \{g_n\}\}$ is the trivial $(n, n, 1, 1)$-SEDF in G. From now on, we concern on nontrivial SEDF's.

The notion of SEDF (and its generalizations) is raised by M. B. Paterson and D. R. Stinson [5] and motivated by its application in communication theory to construct R-optimal regular algebraic manipulation detection codes. Many series of (n, m, k, λ)-SEDF's with $m = 2$ have been constructed in [5, 4, 2, 1]. Paterson and Stinson [5] raised a problem on existence of SEDF with $m \geq 3$. Later, Martin and Stinson [4] proved that there is no (notrivial) SEDF with $m = 3$ and 4. Then Haczunska and Paterson [2] presented more nonexistence results and asked if there exists an SEDF with $m \geq 5$. Very recently, J. Bao et al. [1] presented even more nonexsitence results and conjectured that such SEDF does not exist.

In this note we show an example of SEDF with parameters $(n, m, k, \lambda) = (243, 11, 22, 20)$ which gives an answer of above mentioned problem. More precisely, we will show that the cyclotomic classes $\{C_0, C_1, \ldots, C_{10}\}$ of order 11 in finite field $\mathbb{F}_q$ $(q = 3^5 = 243)$ is a (243, 11, 22, 20)-SEDF in $(\mathbb{F}_q, +)$. Since the example is very concrete, our presentation and computation are down to the earth. Firstly, we determine a primitive element θ of $\mathbb{F}_q$, $\mathbb{F}_q^* = \langle\theta\rangle$. For doing this, we consider the polynomial

$$f(x) = x^5 + x^4 + x^3 + x^2 + 2x + 1 \in \mathbb{F}_3[x].$$

From $g(0) = g(1) = 1$ and $g(2) = g(-1) = -1$ we know that $f(x)$ has no factor $x - a$ $(a \in \mathbb{F}_3)$. Moreover, $f(x)$ is not divided by the quadratic irreducible polynomials $x^2 + 1$ and $x^2 \pm x - 1$ in $\mathbb{F}_3[x]$. Therefore $f(x)$ is irreducible in $\mathbb{F}_3[x]$ and $\mathbb{F}_q = \mathbb{F}_3(\theta)$ where θ is a root of $f(x)$.

Lemma 1. θ is a primitive element of $\mathbb{F}_q$ $(q = 3^5)$.

Proof. Since $\{1, \theta, \theta^2, \theta^3, \theta^4\}$ is a basis of $\mathbb{F}_q$ over $\mathbb{F}_3$, each element α can be expressed uniquely as

$$\alpha = c_0 + c_1\theta + c_2\theta^2 + c_3\theta^3 + c_4\theta^4 \quad (c_i \in \mathbb{F}_3)$$

We identify α as the vector $(c_0, c_1, c_2, c_3, c_4)$ in $\mathbb{F}_3^5$ and denote $\alpha = (c_0c_1c_2c_3c_4)$ briefly. Therefore
$\theta^0 = 1 = (10000), \theta = (01000), \theta^2 = (00100), \theta^3 = (00010), \theta^4 = (00001)$ and $\theta^5 = (21222)$ since $0 = f(\theta) = \theta^5 + \theta^4 + \theta^3 + \theta^2 + 2\theta + 1$.

In general, if $\theta^t = (c_0c_1c_2c_3c_4)$, then

$$\theta^{t+1} = (0c_0c_1c_2c_3) + c_4\theta^5 = (0c_0c_1c_2c_3) + c_4(21222).$$

From this recursive formula, we can get

$$\theta^6 = (11200), \theta^7 = (01120), \theta^8 = (00112), \theta^9 = (12122), \theta^{10} = (10020), \theta^{11} = (01002)$$

and $\theta^{12} = (12211)$. We need to show that the (multiplicative) order of θ is $q-1 = 242 = 2 \cdot 11^2$. For doing this we need to show that $\theta^{22} \neq 1$ and $\theta^{121} \neq 1$. Firstly,

$$\begin{aligned}\theta^{22} &= (\theta^{11})^2 = (\theta + 2\theta^4)^2 = \theta^2 + \theta^5 + \theta^8 = (00100) + (21222) + (00112) \\ &= (21101) \neq 1.\end{aligned}$$

Next

$$\theta^{33} = (\theta^{11})^3 = (\theta + 2\theta^4)^3 = \theta^3 + 2\theta^{12} = (00010) + (21122) = (21102)$$

$$\theta^{99} = (\theta^{33})^3 = (2 + \theta + \theta^2 + 2\theta^4)^3 = 2 + \theta^3 + \theta^6 + 2\theta^{12} = (22002)$$

and

$$\begin{aligned}\theta^{121} &= \theta^{22} \cdot \theta^{99} = (2 + \theta + \theta^2 + \theta^4)(2 + 2\theta + 2\theta^4) = 1 + \theta^2 + 2\theta^3 + \theta^5 + 2\theta^6 + 2\theta^8 \\ &= (20000) = 2 \neq 1.\end{aligned}$$

Therefore θ is a primitive element of $\mathbb{F}_q$ $(q = 3^5)$. □

Next, we need a little knowledge on cyclotomic classes in finite field and cyclotomic numbers. Let $q = p^m$ where $m \geq 1$ and p is a prime, $q - 1 = ef$ $(e \geq 2)$, $\mathbb{F}_q^* = \langle\theta\rangle$. Then $C = \langle\theta^e\rangle$ is the cyclic subgroup of $\mathbb{F}_q^*$ with size $|C| = f$. All cosets $C_\lambda = \theta^\lambda C$ $(0 \leq \lambda \leq e-1)$ of C in $\mathbb{F}_q^*$ are called the cyclotomic classes of order e. The cyclotomic numbers $(i,j)_e$ $(0 \leq i, j \leq e-1)$ over $\mathbb{F}_q$ are defined by

$$(i,j)_e = |(1 + C_i) \cap C_j| = \sharp\{x \in C_i : 1 + x \in C_j\}.$$

The following properties of $(i,j)_e$ can be seen in T. Storer's book [6].

Lemma 2. Let $q = p^m$ where $m \geq 1$ and p is an odd prime, $q - 1 = ef$ $(e \geq 2), C_\lambda$ $(0 \leq \lambda \leq e-1)$ be the cyclotomic classes of order e in $\mathbb{F}_q$, $(i,j)_e$ $(0 \leq i, j \leq e-1)$ be the cyclotomic numbers over $\mathbb{F}_q$. Then

(1) For $r, s, r', s' \in \mathbb{Z}$ and $r \equiv r' (\mathrm{mod} e)$, $s \equiv s' (\mathrm{mod} e)$, we have

$$C_r = C_{r'} \text{ and } (r,s)_e = (r',s')_e.$$

(2)$(i,j)_e = (-i, j-i)_e$ and $(i,j)_e = (pi, pj)_e$ $(0 \leq i, j \leq e-1)$.
(3)$\Delta(C_0, C_0) = f \cdot \{0\} + \sum_{\lambda=0}^{e-1}(e-\lambda, e-\lambda)_e C_\lambda$.
(4)If $2|f$, then $-1 \in C_0$, $-C_\lambda = C_\lambda$, $0 \leq \lambda \leq e-1)$ and $(i,j)_e = (j,i)_e$.

In this paper we need to compute $(i,i)_{11}$ $(0 \leq i \leq 10)$ over $\mathbb{F}_q$ $(q = 3^5)$.

Lemma 3. Let $q = 3^5 = 243, q-1 = ef$ where $e = 11$ and $f = 22$, $(i,j) = (i,j)_{11}$, $(0 \le i,j \le 10)$ be the cyclotomic number of order $e = 11$ over $\mathbb{F}_q$. Then $(0,0) = 1$ and $(i,i) = 2$ for all $1 \le i \le 10$.

Proof. Let $\mathbb{F}_q^* = \langle\theta\rangle$ where θ is the primitive element of $\mathbb{F}_q$ given in Lemma 1. We have the formula

$$\begin{aligned}\Delta(C_0, C_0) &= f\{0\} + \sum_{\lambda=0}^{10}(11-\lambda, 11-\lambda)C_\lambda \text{ (Lemma 2(3))} \\ &= f\{0\} + \sum_{\lambda=0}^{10}(\lambda,\lambda)C_{11-\lambda}\end{aligned}$$

which implies that $f^2 = f + f\sum_{\lambda=0}^{10}(\lambda,\lambda)$. Namely, $\sum_{\lambda=0}^{10}(\lambda,\lambda) = f-1 = 21$. By $(i,i) = (3i,3i)$ (Lemma 2 (2)), we get

$$(1,1) = (3,3) = (9,9) = (5,5) = (4,4), (2,2) = (6,6) = (7,7) = (10,10) = (8,8).$$

Let $A = (0,0), B = (9,9)$ and $C = (7,7)$. Then

$$21 = \sum_{\lambda=0}^{10}(\lambda,\lambda) = A + 5(B+C) \qquad (2)$$

Now we compute the set $C_0 = \langle\theta^{11}\rangle = \{\theta^{11\cdot l} : 0 \le l \le 21\}$ explicitly. From $-1 = \theta^{\frac{q-1}{2}} = \theta^{121} \in C_0$ we know that

$$C_0 = D + (-D),\ D = \{\theta^{11\cdot l} : 0 \le l \le 10\}.$$

We have computed in the proof of Lemma 1 that

$$\theta^{11} = (01002), \theta^{22} = (21101), \theta^{33} = (21102), \theta^{99} = (22002).$$

With the same way of computation we get (by $\theta^{121} = -1$)

$$\theta^{55} = (\theta^{99})^3 = (11112), \theta^{44} = -(\theta^{55})^3 = (12212), \theta^{66} = (\theta^{22})^3 = (10121),$$

$$\theta^{77} = -(\theta^{66})^3 = (12011), \theta^{110} = -(\theta^{77})^3 = (01020), \theta^{88} = (\theta^{110})^3 = (12112).$$

Namely, $C_0 = D + (-D)$ and

$$\begin{aligned}D =& \{\theta^{11\cdot l} : 0 \le l \le 10\} \\ =& \{(10000), (01002), (21101), (21102), (12212), (11112), \\ & (10121), (12011), (12112), (22002), (02010)\}.\end{aligned}$$

From $\pm 1 \in C_0$ and $(-1)-1=1$ we know that $A=(0,0)\geq 1$. From $\theta^{22}=(21101)\in D\subseteq C_0$, $\theta^{33}=(21102)\in D\subseteq C_0$ and $\theta^{33}-\theta^{22}=(00001)=\theta^4$ we know that $\pm\alpha=\pm\theta^{33-4}\in C_7$, $\pm\beta=\pm\theta^{22-4}\in C_7$, $\alpha-\beta=1$, $-\beta-(-\alpha)=1$ and $\alpha=\theta^{29}\neq -\beta=-\theta^{18}$. Therefore $C=(7,7)\geq 2$. Similarly, from $\theta^{44}=(12212)\in D, \theta^{88}=(12112)\in D$ and $\theta^{44}-\theta^{88}=\theta^2$ we know that $\pm\gamma=\pm\theta^{44-2}\in C_9$, $\pm\delta=\pm\theta^{88-2}\in C_9$, $\gamma-\delta=(-\delta)-(-\gamma)=1$ and $\gamma=\theta^{42}\neq -\theta^{86}=-\delta$. Therefore $B=(9,9)\geq 2$. Then by (2) we get

$$21=A+5(B+C)\geq 1+5(2+2)=21,$$

which implies that $A=1$ and $B=C=2$. Namely, $(0,0)=1$ and $(i,i)=2$ for all $1\leq i\leq 10$. This completes the proof of Lemma 3. □

Now we come to the main result.

Theorem 1. Let $q=3^5=243$, $C_\lambda(0\leq\lambda\leq 10)$ be the cyclotomic classes of order 11 of $\mathbb{F}_q$. Then $\{C_0,C_1,\ldots,C_{10}\}$ is an $(n,m,k,\lambda)=(243,11,22,20)$-SEDF in $(\mathbb{F}_q,+)$.

Proof. From $\mathbb{F}_q=\{0\}+\sum_{\lambda=0}^{10}C_\lambda$ we get that for each i $(0\leq i\leq 10)$,

$$\begin{aligned}
\sum_{\substack{j=0\\ j\neq i}}^{10}\Delta(C_i,C_j) &= \Delta(C_i,\sum_{\substack{j=0\\ j\neq i}}^{10}C_j)=\Delta(C_i,\mathbb{F}_q-C_i-\{0\})\\
&= f\mathbb{F}_q-\Delta(C_i,C_i)-C_i \text{ (for any } S\subseteq\mathbb{F}_q,\ \Delta(S,\mathbb{F}_q)=|S|\cdot\mathbb{F}_q)\\
&= f(\mathbb{F}_q-\{0\})-\sum_{\lambda=0}^{10}(i-\lambda,i-\lambda)C_\lambda-C_i\\
&= f(\mathbb{F}_q-\{0\})-(C_i+2\sum_{\substack{\lambda=0\\ \lambda\neq i}}^{10}C_\lambda)-C_i \text{ (by Lemma 3)}\\
&= f(\mathbb{F}_q-\{0\})-2\sum_{\lambda=0}^{10}C_\lambda\\
&= (f-2)(\mathbb{F}_q-\{0\})\\
&= 20(\mathbb{F}_q-\{0\}),
\end{aligned}$$

which means that $\{C_0,C_1,\ldots,C_{10}\}$ is a $(243,11,22,20)$-SEDF in $\mathbb{F}_q$ $(q=3^5)$. □

Remark Our example is not a coincidence. From

$$\Delta(C_i,C_i)=2\sum_{\substack{j=0\\ j\neq i}}^{10}C_j+C_i+f\{0\}=2(\mathbb{F}_q-C_i-\{0\})+C_i+22\{0\}$$

we know that each C_i $(0\leq i\leq 10)$ is an (n,k,λ,μ)-partial difference set (PDS) in $\mathbb{F}_q$ $(q=3^5)$ where $n=q, k=|C_i|=22, \mu=2$ and $\lambda=1=\mu-1$. It is shown ([5],

Theorem 2.4) that if $A_1, \ldots, A_m$ is a partition of an abelian group G and each A_i is a (n, k, λ)-difference set in G, then $\{A_1, \ldots, A_m\}$ is an (n, m, k, λ')-SEDF in G with $\lambda' = k - \lambda$. Similarly we can show that if $A_1, \ldots, A_m$ is a partition of $G - \{0\}$ and each A_i is an (n, k, λ, μ)-PDS in G with $\lambda = \mu - 1$, then $\{A_1, \ldots, A_n\}$ is an (n, m, k, λ')-SEDF in G with $\lambda' = k - \lambda$. In sequential paper we will use the PDS with $\lambda = \mu - 1$ to construct SEDF with $m = 2$, Generalized SEDF and Bounded GSEDF. (For the definition of GSEDF and BGSEDF and their applications we refer to [5]).

On the other hand, it is proved that any nontrivial PDS with $\lambda = \mu - 1$ and $k < \frac{n}{2}$ in a finite abelian group has only two types of parameters $(n, k, \lambda, \mu) = (n, \frac{n-1}{2}, \frac{n-5}{4}, \frac{n-1}{4})$ and $(243, 22, 1, 2)$. The possible SEDF constructed by using these two types of PDS has parameters $m = \frac{n-1}{k} = 2$ and 11. At this moment we still wonder if there exists (n, m, k, λ)-SEDF with $m \geq 5$ and $m \neq 11$.

References

[1] Jingjun Bao, Lijun Ji, Reizhong Wei and Yong Zhang, New existence and nonexistence results for strong external difference families, arXiv: 1612.08385 v1, 26 Dec. 2016.

[2] S. Haczunska and M. B. Paterson, Existence and nonexistence results for strong external families, arXiv: 1611.05621v1. 2016.

[3] S. L. Ma, A survey of partial difference sets, Designs, Codes and Cryptogr. 4(3)(1994), 221-261.

[4] W. Martin and D. R. Stinson, Some nonexistence results for strong external difference families using character theory, arXiv: 1601.06432, 2016.

[5] M. B. Paterson and D. R. Stinson, Combinatorial characterizations of algebraic manipulation detection codes involving generalized difference families, Discrete Math. 339 (2016), 2891-2906.

[6] T. Storer, *Cyclotomy and Difference Sets*, Markham Pub. Co. Chicago, 1967.